Subject Index for *Journal of Ichthyology*, Volume 33, 1993

Abramis brama: **1**, 26–35; **3**, 134–140; **6**, 50–62
Abundance assessment: **8**, 128–133
Acanthochaenus luetkenii: **6**, 85–95
Acipenser baeri: **3**, 9–19; **4**, 15–24; **5**, 66–80; **9**, 85–97
Acipenser gueldenstaedti: **4**, 50–60; **4**, 134–139; **9**, 129–131
Acipenser persicus, Volga River: **1**, 145–146
Acipenser stellatus: **2**, 56–62; **3**, 9–19; **4**, 152–155; **9**, 1–10; **9**, 85–97
Acipenseridae: **3**, 127–133; **8**, 51–65; **9**, 85–97
Adaptive behavior to trawls: **4**, 68–80
Age structure, beluga: **3**, 105–112
Agonidae: **3**, 113–122
Alectrias alectrolophus alectrolophus: **7**, 139–144
Allocareproctus gen. novum: **5**, 99–107
Amur carp:
  Spawning features: **3**, 148–153
  Spawning stock: **3**, 148–153
Anarhichas lupus: **9**, 148–158
Anchovy, Black Sea: **7**, 78–88
Ancient camps, fish remains in: **9**, 134–140
Anguilla anguilla: **7**, 115–123
Astronesthidae: **8**, 111–116
Australia, eastern shores of: **8**, 145–147
Azov basin, introduction of stellate sturgeon: **9**, 1–10

β-phenylethanol: **4**, 50–60; **6**, 118–122
Baikal omul, larval development: **1**, 18–25
Barents Sea: **8**, 18–30
Barguzin River drainage: **3**, 148–153
Behavior changes evoked by β-phenylethanol: **6**, 118–122
Behavior differences: **9**, 141–147
Behavior in turbulent flow: **5**, 45–55
Beluga: **9**, 132–133
Beluga, Caspian Sea: **9**, 72–84
Beluga migration into Volga: **3**, 105–112
Bering Sea, walleye pollock: **2**, 63–76
Beryciformes: **3**, 30–49
Biostatistical methods: **8**, 96–110
Black carp: **3**, 66–75
Black Sea: **7**, 100–114
Black Sea basin: **8**, 122–127
Blenny, developing eggs of: **1**, 149–152
Blood leukocytes in population screening: **2**, 151–158
Blue whiting: **2**, 77–86; **5**, 26–34
  Biology: **5**, 26–34
  Distribution: **5**, 26–34
  West European basin: **2**, 77–86
Bol'shaya River: **5**, 150–154
Bol'shaya River Estuary: **6**, 153–158; **9**, 11–21
Bomolochidae: **6**, 63–67
Bomolochus myctophi sp. n.: **6**, 63–67
Brachydanio rerio: **6**, 118–122
Brachymystax lenok: **8**, 148–156
Brackish water: **7**, 1–11
Bream: **4**, 68–80; **6**, 50–62
  Biological parameters: **3**, 134–140
  Fishery dynamics: **3**, 134–140
Brown trout: **2**, 118–124
Burbot morphology: **4**, 140–144

Capelin: **8**, 18–30
Carassius carassius: **2**, 151–158
Carbohydrate transport:
  In bream: **1**, 26–35
  In carp: **1**, 26–35
Carp population:
  Ecological characteristics: **6**, 141–147
  Genetic characteristics: **6**, 141–147
Cartilaginous fishes: **4**, 111–126
Caspian Sea: **9**, 1–10
Caudal region: **9**, 132–133
Centrolophidae: **2**, 125–139; **4**, 1–14

Channel catfish: **1**, 85–92; **2**, 87–103; **5**, 154–158
Early development: **2**, 87–103
Channel groupings: **9**, 22–30
Chara River round whitefish: **2**, 37–47
Chars, freshwater: **3**, 57–65
Chemical signals, food: **8**, 51–65
Chemical stimuli, natural: **4**, 50–60
Chemoreceptor organs: **5**, 143–149
Chemosensory-organ development: **3**, 9–19
Cherry salmon: **6**, 148–152
Chondrichthyes: **4**, 111–126
Choridactylus, new species: **5**, 136–142
Chu River basin, percids: **3**, 76–85
Chum salmon: **2**, 104–116; **4**, 127–133
Cisco: **7**, 1–11
Clupea pallasi marisalbi: **8**, 75–86; **8**, 87–95
Cobitidae (Balitoridae): **4**, 145–147
Cod, abundance of Eastern Baltic: **8**, 96–110
Coelorinchus fasciatus: **1**, 71–84
Coho salmon: **7**, 145–146
Cohort analysis: **1**, 63–70
Comephoridae, Lake Baikal: **4**, 61–67
Commercial fishing as selection factor: **9**, 141–147
Commercial return, stellate sturgeon: **2**, 56–62
Commercially important fishes: **7**, 100–114
Conditioned reflex in fishes: **8**, 40–50
Coregonus autumnalis migratorius: **1**, 18–25
Coregonus peled: **4**, 148–151
Coregonus sardinella: **7**, 1–11
Cottoid fish: **4**, 61–67
Cottoidei: **4**, 61–67
Craniological analysis: **5**, 108–123; **6**, 96–106
Crucian carp: **2**, 151–158
Ctenopharyngodon idella: **3**, 66–75
Cyprinidae: **9**, 98–116
Cyprinus carpio haematopterus: **3**, 148–153
Cyprinus carpio: **1**, 26–35; **1**, 54–62; **4**, 81–92

Dace: **5**, 45–55; **9**, 98–116
Defensive behavior, bream: **4**, 68–80
Deformed vertebral column: **9**, 132–133
Diaphus benthopelagic species: **3**, 20–29
Diaphus coeruleus: **8**, 140–144
Digestive tract:
Morphogenesis: **4**, 111–126
Morphology: **4**, 111–126
Digestive-enzyme activity: **9**, 48–56
Distribution dynamics, seasonal spatial: **2**, 63–76
Distribution of pink salmon: **4**, 38–49
Double-fan model: **5**, 35–44

East China Sea: **8**, 117–121
East Siberian Sea: **2**, 26–36
Ecological studies: **2**, 151–158
Ecology of *Diaphus coeruleus*: **8**, 140–144
Ectoparasites: **6**, 50–62
Ectopic locations of sex cells, sturgeons: **2**, 125–139
Egg membrane formation: **1**, 85–92
Egg membrane mechanical properties: **5**, 124–129
Elassodiscus, validity of genus: **6**, 68–84
Embryogenesis period: **5**, 124–129
Embryonal movement: **5**, 124–129
Embryonic gonad, formation of: **7**, 115–123
Endangered (rare) fishes: **1**, 109–127
Energetics of juvenile fish: **3**, 1–8
Energy exchange dynamics: **1**, 18–25
Engraulis encrasicholus: **7**, 78–88
Equilibrium in fish: **5**, 45–55
Ethologometry: **8**, 31–39
Eupogonesthes xenicus: **8**, 111–116
European eel: **7**, 115–123
Exocoetidae: **5**, 12–25
Experimental modeling: **1**, 63–70

Feeding, daily dynamics of: **7**, 78–88
Feeding specializations: **9**, 122–128
Fish health in aquaculture: **1**, 44–53
Fish population:
Distribution: **7**, 100–114
Estimates: **7**, 100–114
Fish productivity: **5**, 1–11
Fishing gear, avoidance of: **8**, 40–50
Flounder: **7**, 124–138

Flying fishes: **5**, 12–25
Fossil Perciformes, Caucasian: **3**, 50–56
Freshwater fishes, list: **4**, 99–110

Gadidae: **1**, 101–108
Gadiformes: **8**, 117–121
Gadus morhua callarias: **8**, 96–110
Gadus morhua morhua: **9**, 148–158
Genetic variability, brown trout: **2**, 118–124
Gilbertidia, new species of: **6**, 130–133
Gobiidae: **5**, 108–123; **8**, 122–127
  Black Sea basin: **1**, 137–143
  Caspian Sea basin: **1**, 137–143
  Mediterranean region: **2**, 1–16
  Ponto-Caspian region: **2**, 1–16
Goby: **5**, 108–123; **6**, 96–106; **8**, 122–127
  Karyotypes: **1**, 137–143
Gonad maturity stages: **4**, 1–14
Gonadogenesis, disruption of: **3**, 127–133
Gonads, prespawning state: **7**, 12–22
Grass carp: **3**, 66–75
Gray mullet: **7**, 152–158
Gray Notothenia: **9**, 57–71
Grayling:
  Arctic: **7**, 89–99
  Biological characteristics of, in northeastern Asia: **7**, 89–99
  Oocyte structure: **1**, 9–17
Greenlings: **4**, 25–37
  Growth, physiological aspects of: **6**, 123–129
  One-finned: **6**, 123–129
Grenadier:
  Discovery of rare species: **8**, 117–121
  Reproductive biology: **1**, 71–84
Gulf of Aden: **5**, 136–142
Gustatory (stimulating) substances: **9**, 85–97

Hake, Cape: **8**, 128–133; **9**, 117–121
Hatchery-reared chum salmon: **1**, 36–43
Hemigrammus caudovittatus: **8**, 40–50
Herring:
  White Sea: **8**, 75–86; **8**, 87–95
  White Sea, spatial dynamics of year classes: **8**, 87–95
Heterochromatin: **6**, 42–49
Homeostasis (salt) in sturgeon: **1**, 92–100
Homeostasis (water) in sturgeon: **1**, 92–100
Hormonal stimulation: **4**, 127–133
Huso huso: **3**, 105–112; **9**, 72–84; **9**, 132–133
Hybrids (reciprocal): **3**, 66–75
Hymenocephalus papyraceus: **8**, 117–121
Hypomesus olidus: **9**, 11–21

Ichthyofauna: **6**, 136–140
  Central Asia: **4**, 99–110
Ichthyonekton: **5**, 1–11
Ichthyophthirius multifilis: **4**, 81–92
Ichthyoplankton: **5**, 1–11
  On seamounts: **2**, 17–25
  Western Indian Ocean: **2**, 17–25
Ictalurus punctatus: **1**, 85–92; **2**, 87–103; **5**, 154–158
Idiolychnus benthopelagic species: **3**, 20–29
Indian butterfish: **2**, 125–139
Indian Ocean:
  Eastern: **8**, 111–116
  West, recent fish collections: **1**, 128–136
Irgiz-Turagaiskii basin: **6**, 136–140
Iturup Island coastal waters: **6**, 1–14

Jack mackerel, Peruvian: **3**, 86–104
Japan Trench: **9**, 31–39
Jaw-pharyngeal apparatus: **9**, 122–128
Juvenile chum salmon: **1**, 36–43; **4**, 127–133; **6**, 15–22
Juvenile Pacific salmon: **6**, 15–22
Juvenile salmon: **6**, 1–14

Kamchatka River basin: **2**, 104–116; **3**, 57–65
Karyology of Gobiidae: **2**, 1–16
Karyotype (unique): **6**, 42–49
Karyotypes: **8**, 122–127
Kazakhstan carp: **1**, 54–62
  Biochemical characters: **1**, 54–62
  External characters: **1**, 54–62
Kentrocapros rosapinto: **6**, 134–135
Kolyma River: **5**, 66–80
Kuril Islands: **6**, 130–133; **7**, 139–144
Kuril-Kamchatka Trench: **5**, 130–135

Kyushyu-Palau submarine ridge: **3**, 30–49

Lactate dehydrogenase reaction: **1**, 101–108
Laemonema longipes: **6**, 107–117
  Distribution of: **6**, 107–117
  Migration of: **6**, 107–117
Lantern fish, spiny: **6**, 63–67
Lateral-line organs: **5**, 45–55
Learning-ability differences: **9**, 141–147
Lena River: **7**, 1–11
Lena Tablemount: **9**, 57–71
Leuciscus danilewski: **9**, 98–116
Leuciscus leuciscus: **5**, 45–55
Leuciscus subgenus: **9**, 98–116
Liparidae: **5**, 81–98; **5**, 99–107; **6**, 68–84; **9**, 31–39
Littoral fish: **7**, 139–144
Littoral groupings: **9**, 22–30
Liza ramada: **7**, 152–158
Loach, distribution: **4**, 145–147
Lota lota: **4**, 140–144
Lycenchelys makushok sp. nova: **5**, 130–135

M. carinatus distribution: **1**, 71–84
M. whitsoni: **1**, 71–84
Macrouridae: **1**, 71–84; **8**, 117–121
Macrourus carinatus: **1**, 71–84
Maldive Islands: **6**, 134–135
Mallotus villosus: **8**, 18–30
Maturation cycles: **9**, 72–84
Merluccius capensis: **8**, 128–133; **9**, 117–121
Mesobenthic fish, new: **8**, 111–116
Mesopelagic fishes: **8**, 1–17
Metabolic rate, effect of feeding on: **9**, 148–158
Metabolism: **3**, 1–8
Methyltestosterone in carp: **1**, 147–148
Micromesistius poutassou: **2**, 77–86; **5**, 26–34
Micropyle number variability in eggs: **1**, 145–146
Micropyles in eggs: **9**, 129–131
Mid-Atlantic Ridge: **5**, 26–34
Migration, anadromous, of salmon: **7**, 63–77
Minnow, common: **9**, 22–30
Minous, new species: **5**, 136–142
Moridae: **6**, 107–117
Morph formation: **3**, 57–65
Muscle degeneration in sturgeon: **1**, 92–100
Muscle degeneration pathology: **4**, 134–139
Myctophidae: **3**, 20–29; **4**, 93–98; **8**, 134–139
Myctophum ovcharovi: **4**, 93–98
Myctophum spinosum: **6**, 63–67
Mylopharyngodon piceus: **3**, 66–75

Neogobius cephalargoides: **5**, 108–123
Neogobius platyrostris: **5**, 108–123
Naimin Nuur Lakes (Mongolia): **4**, 148–151
New species:
  Centriscidae, Syngnathiformes: **3**, 123–126
  Genus *Elassodiscus*: **6**, 68–84
  Genus *Palaeorhynchus*: **3**, 50–56
  Gilbertidia: **6**, 130–133
  Myctophum asperum: **4**, 93–98
  Southeastern Pacific Ocean: **3**, 20–29
Noemacheilus: **4**, 145–147
Northern-latitude fishes: **9**, 48–56
Northwestern Pacific: **6**, 107–117
Notothenia squamifrons squamifrons: **9**, 57–71
Nototheniidae: **1**, 71–84; **7**, 23–47
Notothenioidei: **7**, 23–47

Ob Tablemount: **9**, 57–71
Olfactory sensitivity: **8**, 51–65
Oncorhynchus gorbuscha: **4**, 38–49; **6**, 148–152
Oncorhynchus keta: **1**, 36–43; **2**, 104–116; **4**, 127–133; **6**, 15–22
Oncorhynchus kisutch: **6**, 15–22; **7**, 145–146
Oncorhynchus masou: **6**, 148–152
Oncorhynchus mykiss: **8**, 66–74
Oncorhynchus nerka: **5**, 35–44
Ontogeny:
  Early, effect of human activity on: **6**, 23–41
  Sturgeon: **8**, 51–65
Oocyte degeneration: **1**, 1–8
Oogenesis, periods of: **2**, 125–139

Osmerus mordax dentex: **5**, 150–154
Osmoregulatory capability: **1**, 36–43
Ostraciontidae: **6**, 134–135
Ovarian development, Indian butterfish: **2**, 125–139
Oxygen consumption: **6**, 15–22

Pacific salmon: **5**, 56–65
  Freshwater residence: **5**, 56–65
Pacific sardine: **1**, 1–8
Paddlefish: **5**, 143–149
Palaeorhynchidae: **3**, 50–56
Palaeorhynchus parini: **3**, 50–56
Parasites: **6**, 63–67
Patagonotothen guentheri shagensis: **1**, 71–84
Pechora River basin: **9**, 48–56
Peled:
  Age composition: **4**, 148–151
  Growth: **4**, 148–151
Percids: **3**, 76–85
Perciformes: **3**, 50–56; **5**, 130–135
Petrel: **7**, 145–146
Petroleum fractions, effect of: **1**, 149–152
Phenetic diversity: **9**, 98–116
Photophores, fish-skin: **8**, 1–17
Phoxinus phoxinus: **9**, 22–30
Phylogeny of Nototheniidae: **7**, 23–47
Physiological energetics: **5**, 35–44
Phytophilic fishes: **7**, 12–22
Pink salmon: **4**, 38–49; **6**, 148–152
Pleurogrammus azonus: **6**, 123–129
Pleurogrammus monopterygius: **4**, 25–37
Pleuronectes platessa: **9**, 148–158
Pleuronectidae: **7**, 124–138
Pleuronectiform fishes: **9**, 122–128
Poecilopsetta: **7**, 124–138
Pollution, water: **6**, 50–62
Polymixia, new species: **3**, 30–49
Polymixia busakhini: **8**, 145–147
Polymixiidae: **3**, 30–49
Polyodon spathula: **5**, 143–149
Polyodontidae: **5**, 143–149
Pond smelt, diet of: **9**, 11–21
Ponticola Iljin, subgenus: **5**, 108–123; **6**, 96–106
Population dynamics:
  Beluga: **3**, 105–112
  Herring: **2**, 48–55
Population modeling: **8**, 75–86
Population organization, Pacific salmon: **5**, 56–65
Poronay River, Sakhalin: **4**, 140–144
Preservation strategies: **1**, 109–127
Previtelligenous oocytes: **1**, 1–8
Primorye: **6**, 123–129
Prosopium cylandraceum: **2**, 37–47
Protective adaptation: **8**, 1–17
Proval Cove: **8**, 128–133
Psednos: **5**, 81–98
Psenopsis cyanea: **2**, 125–139; **4**, 1–14
Psetta maeotica: **5**, 124–129
Pseudoliparis, new species description: **9**, 31–39
Psychrolutidae: **6**, 130–133
Puntius conchonius: **8**, 40–50

Quaternary freshwater fishes: **2**, 26–36

Radioactive waters: **6**, 141–147
Rainbow trout: **8**, 66–74
Reproduction, greenlings: **4**, 25–37
Reproductive isolates: **7**, 12–22
Reproductive system as bioindicator: **4**, 15–24
Roach, juvenile: **9**, 22–30
Round whitefish biological data: **2**, 37–47
Rutilus rutilus: **9**, 22–30
Rybinsk Reservoir: **3**, 134–140; **7**, 12–22

Salinity fluctuations: **3**, 1–8
Salinity, effect of: **6**, 15–22
Salmo gairdneri: **8**, 66–74
Salmo mykiss: **1**, 9–17
Salmo trutta: **2**, 118–124
Salmon:
  Anadromous migration of: **7**, 63–77
  Distribution in Sakhalin–Kuril waters: **7**, 63–77
  Distribution of in Bering Sea: **7**, 48–62
  Migration of in Bering Sea and Pacific waters: **7**, 48–62

Salmonid oocyte structure: **1**, 9–17
Salmonids, phylogeny of: **8**, 148–156
Salmonoidei suborder: **1**, 9–17
Salvelinus: **3**, 57–65
Salvelinus alpinus: **8**, 148–156
Salvelinus namaycush: **8**, 148–156
Salvethymus svetovidovi: **6**, 42–49
Samarga River: **6**, 148–152
Sardine, Far East: **7**, 147–151
Sardinops sagax melanosticta: **1**, 1–8; **7**, 147–151
Scorpaeniformes: **5**, 81–98; **5**, 99–107; **5**, 136–142; **6**, 68–84; **6**, 130–133; **9**, 31–39
Sea of Azov: **6**, 96–106
Sea of Okhotsk: **4**, 145–147; **6**, 68–84; **9**, 134–140
Sea poachers: **3**, 113–122
Seamount communities: **2**, 17–25
Seasonality of growth: **7**, 1–11; **7**, 147–151
Selection significance of biochemical characters: **1**, 54–62
Selection significance of external characters: **1**, 54–62
Sex cells, annual pool: **4**, 1–14
Sex cycle, Centrolophidae: **4**, 1–14
Sexual cycle: **1**, 1–8
Sexual dimorphism: **3**, 113–122
Shchugor River: **9**, 48–56
Shikotan Island: **7**, 139–144
Shrimpfishes, fossil: **3**, 123–126
Siberian sturgeon: **3**, 9–19; **4**, 15–24; **5**, 66–80; **9**, 85–97
Smelt larvae diet: **5**, 150–154
Smelt: **5**, 150–154
Snailfishes, new genus: **5**, 99–107
Sockeye salmon: **5**, 35–44
Southeastern India coastal waters: **5**, 12–25
Spatial subdivision of capelin: **8**, 18–30
Spawning, blue whiting: **2**, 77–86
Spawning, environmental influences of: **2**, 104–116
Spawning sites, chum salmon: **2**, 104–116
Spawning stock, capelin: **8**, 18–30
Sperm production: **7**, 152–158
Spermatogenesis, gray mullet: **7**, 152–158
Starry flounder, size–age structure: **6**, 153–158
Stellate sturgeon: **9**, 1–10; **9**, 85–97
  Volga: **4**, 152–155
  Survival rate: **2**, 56–62
Stephanoberycidae: **6**, 85–95
Stichaeidae: **7**, 139–144
Stichaeopsis nana: **7**, 139–144
Stress, aspects of: **8**, 66–74
Sturgeon:
  Caspian Sea: **2**, 140–150; **3**, 127–133
  Russian: **4**, 50–60; **4**, 134–139; **9**, 129–131
Subtropical waters: **5**, 1–11
Supporting-elements morphology: **9**, 122–128
Swordfish, spawning, Black Sea: **1**, 144
Synanceiidae: **5**, 136–142
Syngnathiformes, fossil: **3**, 123–126

Taxonomic relationships: **8**, 122–127
Taxonomy of Gobiidae: **2**, 1–16
Temporal subdivision of capelin: **8**, 18–30
Theragra chalcogramma: **2**, 63–76; **3**, 141–147
Thymallus arcticus: **1**, 9–17; **8**, 148–156
Trachurus symmetricus murphyi: **3**, 86–104
Trap-net model: **9**, 141–147
Trawl surveys: **9**, 117–121
Trophic interrelationships: **6**, 1–14
Tuna, feeding habits: **9**, 40–47
Turbot, Black Sea: **5**, 124–129

Ultra-abyssal genus: **9**, 31–39
Underyearling carp: **4**, 81–92
Unfertilized eggs: **5**, 154–158
Upper Eocene bony fishes: **3**, 50–56
Upper Eocene shrimpfishes: **3**, 123–126
Upwelling indicators, local: **9**, 40–47

Validity of genus *Psednos*: **5**, 81–98
Viability of chum salmon progeny: **2**, 104–116
Viability of eggs: **5**, 154–158
Visceral skeleton of Nototheniidae: **7**, 23–47
Vitelligenous oocytes: **1**, 1–8

Volga River: **9**, 129–131
Volga sturgeon: **6**, 23–41
Vomer, development of: **8**, 148–156

Walleye pollock: **2**, 63–76
  Year class abundance: **3**, 141–147
West Bering Sea: **3**, 141–147
White Sea herring: **2**, 48–55

Xenopelagic deepwater fishes: **9**, 40–47
Xiphias gladius, Black Sea: **1**, 144

Year-class strength assessment: **9**, 117–121

Zoarcidae: **5**, 130–135
Zoogeographic analysis: **4**, 99–110